

Calabi-Yau Black Holes and Enhancement of Supersymmetry in Five Dimensions

A. H. Chamseddine and W. A. Sabra ¹

*Center for Advanced Mathematical Sciences (CAMS),
and
Physics Department, American University of Beirut, Lebanon*

ABSTRACT

BPS electric and magnetic black hole solutions which break half of supersymmetry in the theory of $N = 2$ five-dimensional supergravity are discussed. For models which arise as compactifications of M-theory on a Calabi-Yau manifold, these solutions correspond, respectively, to the two and five branes wrapping around the homology cycles of the Calabi-Yau compact space. The electric solutions are reviewed and the magnetic solutions are constructed. The near-horizon physics of these solutions is examined and in particular the phenomenon of the enhancement of supersymmetry. The solutions for the supersymmetric Killing spinor of the near horizon geometry, identified as $AdS_3 \times S^2$ and $AdS_2 \times S^3$ are also given.

¹ e-mail: chams@aub.edu.lb, ws00@aub.edu.lb

1 Introduction

Considerable progress has been made in the study of BPS black hole states of the low-energy effective actions of compactified string and M -theory². This was mainly motivated by the important role that these states play in the understanding of the non-perturbative structure behind string theory. Toroidal compactifications of string theory give rise to vacua with $N = 4$ and $N = 8$ supersymmetry and BPS states of these models are restricted by the large supersymmetry. Corrections to these solutions and their entropies can only arise from higher loops, as the lowest order corrections are known to vanish. Four and five dimensional $N = 2$ supergravity models coupled to vector and hyper-multiplets can arise, for example, from type II string and M -theory compactified on a Calabi-Yau threefold. An important step in the study of $N = 2$ black holes was the realisation that their entropy is given in terms of the extremum of the central charge and that the scalar fields of the theory take fixed values at the horizon independent of their boundary values at spatial infinity [1].

The perturbative and nonperturbative corrections for the $N = 2$ supergravity models seem to make the study of their BPS black hole solutions more intricate. However, the rich geometric structure of these theories [3, 4] considerably simplifies the analysis of their black hole solutions. For instance, the black hole metric in four dimensions can be expressed in terms of symplectic invariant quantities where the symplectic sections satisfy algebraic constraints involving a set of constrained harmonic functions [5]. In a similar way to Einstein-Maxwell theory [6], different types of solutions can be obtained depending on the choice of harmonic functions. The solutions also depend on the choice of the prepotential which defines the theory. In five dimensions, the black hole solutions [7] can be expressed in terms of the rescaled cubic homogeneous prepotential which defines very special geometry [4].

The BPS solutions can be regarded as solitons interpolating between two vacua: Minkowski flat space at infinity and $AdS_{p+2} \times S^{d-p-2}$ near the horizon [8, 9, 10], where p is the dimensionality of the solution. Anti-de Sitter spaces and spheres are known to admit Killing spinors which we refer to as geometric Killing spinors. At a generic point in space-time, the BPS solution breaks half of supersymmetry. However, if at the near horizon the supersymmetric Killing spinor, defined by the zero mode of the gravitino supersymmetry transformation, can be identified with the geometric spinor, then supersymmetry is enhanced. This restoration of supersymmetry have been discussed in the literature for the cases where the near horizon geometry was identified with the spaces $AdS_2 \times S^2$ [8, 9], $AdS_5 \times S^5$, $AdS_3 \times S^7$, and $AdS_7 \times S^3$ [10]. In five dimensional supergravity models with vector multiplets, our concern in this paper, the restoration of supersymmetry of a double-extreme black hole solutions [11] near the horizon in four and five dimensions has been discussed in [1, 12] and also for the rotating case in [13]. The analysis of the pure five dimensional supergravity case (without vector multiplets) was

²for reviews see for example [2]

performed in [14].

General BPS black hole solutions which break half of supersymmetry of $d = 5$, $N = 2$ supergravity theory with an arbitrary number of vector supermultiplets, were obtained in [7]. There it was also demonstrated that the entropy of these black holes is given in terms of the extremised central charge. The near horizon geometry of these solutions is the same as that of the double-extreme solutions in accordance with the attractor behaviour. The main result of this paper is the analysis of general magnetic-string solutions³, which in compactified M-theory on a Calabi-Yau manifold, correspond to five branes wrapped around the four cycles of the Calabi-Yau compact space. The enhancement of supersymmetry at the near horizon of these solutions is shown and we derive the expression of the Killing spinor of the maximally supersymmetric horizon. The analysis of [12] is also completed by obtaining the form of the Killing spinor for the near horizon geometry for the electric BPS solution given by $AdS_2 \times S_3$.

The bosonic action of five dimensional $N = 2$ supergravity coupled to vector multiplets is given by

$$e^{-1}\mathcal{L} = -\frac{1}{2}R - \frac{1}{4}G_{IJ}F_{\mu\nu}^I F^{\mu\nu J} - \frac{1}{2}g_{ij}\partial_\mu\phi^i\partial^\mu\phi^j + \frac{e^{-1}}{48}\epsilon^{\mu\nu\rho\sigma\lambda}C_{IJK}F_{\mu\nu}^I F_{\rho\sigma}^J A_\lambda^K. \quad (1)$$

The theory of five dimensional $N = 2$ supergravity with vector supermultiplets was considered in [17]. The compactification of $N = 1$, $D = 11$ supergravity [16], i. e., the low energy limit of M-theory, down to five dimensions on Calabi-Yau 3-folds (CY_3), results in a $d = 5$, $N = 2$ supergravity coupled to abelian vector supermultiplets based on the structure of very special geometry [4]. The five dimensional theory contains the gravity multiplet, $h_{(1,1)} - 1$ vector multiplets and $h_{(2,1)} + 1$ hypermultiplets, $(h_{(1,1)}, h_{(2,1)})$ are the Hodge numbers of CY_3 . The $(h_{(1,1)} - 1)$ -dimensional space of scalar components of the abelian vector supermultiplets coupled to supergravity can be regarded as a hypersurface of a $h_{(1,1)}$ -dimensional manifold whose coordinates $X^I(\phi)$ are in correspondence with the vector bosons (including the graviphoton). In five dimensions, an $N = 2$ vector supermultiplet has a single scalar field and therefore the scalar manifold is real. The defining equation of the hypersurface is

$$\mathcal{V}(X) = 1 \quad (2)$$

and the prepotential $\mathcal{V}$ is a homogeneous cubic polynomial in the coordinates $X^I(\phi)$

$$\mathcal{V}(X) = \frac{1}{6} C_{IJK} X^I X^J X^K = X^I X_I = 1, \quad I, J, K = 1, \dots, h_{(1,1)} \quad (3)$$

where C_{IJK} are the topological intersection numbers and X_I are the “dual” special coordinates (the four cycles).

³ a magnetic string solution in five dimensions was obtained as a compactification of an intersection of 3 M-5-branes and a boost along the common string in [15]

The corresponding vector and scalar metrics in (1) are completely encoded in the function $\mathcal{V}(X)$

$$\begin{aligned} G_{IJ} &= -\frac{1}{2} \frac{\partial}{\partial X^I} \frac{\partial}{\partial X^J} \ln \mathcal{V}(X)|_{\mathcal{V}=1}, \\ g_{ij} &= G_{IJ} \frac{\partial}{\partial \phi^i} X^I \frac{\partial}{\partial \phi^j} X^J|_{\mathcal{V}=1}. \end{aligned} \quad (4)$$

2 Electric BPS Black Holes

In the compactified M-theory on a Calabi-Yau manifold, electrically point and magnetically charged string-like BPS states correspond to the two- and five-branes of M-theory wrapped around the two- and four-cycles of the Calabi-Yau space. Though the details of the low-energy Lagrangian depend very much on the compactified Calabi-Yau space, the analysis of the BPS solutions can be considerably simplified by the rich geometric structure based on “very special geometry” [4] underlying the $N = 2$ five-dimensional theories with vector supermultiplets. Electrically charged BPS solutions of the five dimensional theory have been analyzed in [7] and will be reviewed in this section.

The supersymmetry transformation laws for the fermi fields in a bosonic background are given by [7, 17]

$$\begin{aligned} \delta\psi_\mu &= \left(\mathcal{D}_\mu + \frac{i}{8} X_I (\Gamma_\mu^{\nu\rho} - 4\delta_\mu^\nu \Gamma^\rho) F_{\nu\rho}^I \right) \epsilon, \\ \delta\lambda_i &= \left(\frac{3}{8} \partial_i X_I \Gamma^{\mu\nu} F_{\mu\nu}^I - \frac{i}{2} g_{ij} \Gamma^\mu \partial_\mu \phi^j \right) \epsilon, \end{aligned} \quad (5)$$

where ϵ is the supersymmetry parameter and $\mathcal{D}_\mu$ the covariant derivative.⁴ The metric for the BPS electric black hole solutions can be brought to the form [7]

$$ds^2 = -e^{-4U} (dt + w_m dx^m)^2 + e^{2U} (d\vec{x})^2. \quad (6)$$

The vanishing of the supersymmetry transformations of the gravitino and gauginos, for the choice $\Gamma^0 \epsilon = -i\epsilon$, imposes the following relations for the supersymmetry variation parameter, U , w_m , the scalar fields and the gauge fields

$$\begin{aligned} \partial_t \epsilon &= 0, \\ (\partial_m + \partial_m U) \epsilon &= 0, \\ (X_I F_{mn}^I)^- &= (\partial_m Q_n - \partial_n Q_m)^-, \\ (X_I F_{tm}^I) &= -\partial_m e^{-2U}, \\ 2(\partial_m U w_n - \partial_n U w_m) - \frac{1}{2}(\partial_m w_n - \partial_n w_m) + e^{2U} (X_I F_{mq}^I) &= 0. \end{aligned}$$

⁴ we use the metric $\eta^{ab} = (-, +, +, +, +)$, $\{\Gamma^a, \Gamma^b\} = 2\eta^{ab}$, $\mathcal{D}_\mu = \partial_\mu + \frac{1}{4}\omega_{\mu ab}\Gamma^{ab}$, $\omega_{\mu ab}$ is the spin connection, and Γ^ν are Dirac matrices and $\Gamma^{a_1 a_2 \dots a_n} = \frac{1}{n!} \Gamma^{[a_1} \Gamma^{a_2} \dots \Gamma^{a_n]}$.

$$\begin{aligned}
-\frac{1}{2}e^{2U}(X_I F_{mq}^I)^* - (\partial_m U w_n - \partial_n U w_m)^* &= 0, \\
\frac{3}{2}e^{-U}\partial_m X_I \partial_i X^I - G_{IJ}e^U \partial_i X^I F_{tm}^J &= 0, \\
\left(G_{IJ}\partial_i X^I F_{mn}^J + \frac{3}{2}e^{-2U}(\partial_m X_I w_n - \partial_n X_I w_m)\right)^- &= 0.
\end{aligned} \tag{7}$$

where $Q_n \equiv e^{-2U}w_n$ and $F_{mn}^- = F_{mn} - {}^*F_{mn}$. The above conditions are satisfied for

$$\begin{aligned}
F_{mn}^I &= \partial_m(X^I Q_n) - \partial_n(X^I Q_m), \\
F_{tm}^I &= -\partial_m(e^{-2U}X^I), \\
(\partial_m w_n - \partial_n w_m)^- &= 0 \\
\epsilon &= e^{-U}\epsilon_0.
\end{aligned} \tag{8}$$

where ϵ_0 is a constant spinor satisfying $\Gamma^0 \epsilon = -i\epsilon$.

The various physical variables can be fixed in terms of space-time functions, by solving the equations of motion for the gauge fields⁵. From the Lagrangian (1), we derive the following equation of motion for the gauge fields,

$$\partial_\nu(eG_{IJ}g^{\mu\rho}g^{\nu\sigma}F_{\rho\sigma}^J) = \frac{1}{16}C_{IJK}\epsilon^{\mu\nu\rho\sigma k}F_{\nu\rho}^J F_{\sigma k}^K. \tag{9}$$

Using (8) and after some manipulations one obtains the solution

$$e^{2U}X_I = \frac{1}{3}H_I, \tag{10}$$

where H_I are harmonic functions, $H_I = h_I + \frac{q_I}{r^2}$, h_I are constants and q_I are electric charges.

It is convenient to express the solution in terms of the geometry of the internal space, i. e., in terms of the cubic polynomial $\mathcal{V}$. Define the rescaled coordinates

$$Y_I = e^{2U}X_I, \quad Y^I = e^U X^I, \tag{11}$$

then the underlying very special geometry implies that

$$e^{3U} = \mathcal{V}(Y) = \frac{1}{3}C_{IJK}Y^I Y^J Y^K, \tag{12}$$

and thus the electric black hole solution metric has the form

$$\begin{aligned}
ds^2 &= -\mathcal{V}^{-4/3}(Y)(dt + \omega_m dx^m)^2 + \mathcal{V}^{2/3}(Y)(d\vec{x})^2, \\
Y_I &= \frac{1}{3}H_I, \\
F_{mn}^I &= \partial_m(\mathcal{V}^{-1}Y^I w_n) - \partial_n(\mathcal{V}^{-1}Y^I w_m), \\
F_{tm}^I &= -\partial_m(\mathcal{V}^{-1}Y^I).
\end{aligned} \tag{13}$$

⁵ notice that the Bianchi identities are trivially satisfied

Black holes with rotational symmetry can be constructed using the above general solution. Changing to polar coordinates defined by, $x^1 + ix^2 = r \sin \theta e^{i\phi}$, $x^3 + ix^4 = r \cos \theta e^{i\psi}$, and specializing to solutions with rotational symmetry in the two orthogonal planes, i. e., $w_\phi = w_\phi(r, \theta)$, $w_\psi = w_\psi(r, \theta)$, $w_r = \omega_\theta = 0$, then the self-duality condition of the field strength of w_m gives for a decaying solution, $w_\phi = -\frac{\alpha}{r^2} \sin^2 \theta$, $w_\psi = \frac{\alpha}{r^2} \cos^2 \theta$. Therefore, the general form of the metric solution with rotational symmetry in the two orthogonal planes is given by

$$ds^2 = -\mathcal{V}^{-4/3}(Y) \left(dt - \frac{\alpha \sin^2 \theta}{r^2} d\phi + \frac{\alpha \cos^2 \theta}{r^2} d\psi \right)^2 + \mathcal{V}^{2/3} (dr^2 + r^2 d\Omega_3^2) \quad (14)$$

where

$$d\Omega_3^2 = (d\theta^2 + \sin^2 \theta d\phi^2 + \cos^2 \theta d\psi^2). \quad (15)$$

It was demonstrated in [12] that for double-extreme black holes in five dimensions supersymmetry is enhanced at the near horizon where the metric takes the form

$$ds^2 = -\left(\frac{2\hat{r}}{r_0}\right)^2 dt^2 + \left(\frac{2\hat{r}}{r_0}\right)^{-2} d\hat{r}^2 + r_0^2 d\Omega_3^2, \quad (16)$$

with the geometry $AdS_2 \times S^3$. Also these results were extended to the rotating case where similar conclusions hold [13]. In reality one can obtain the stronger result that for a general electric BPS black hole in five dimensions with vector supermultiplets, supersymmetry is restored at the near-horizon. This is because both extreme and double-extreme black holes have the same form of metric at the near horizon. This can be seen by examining the general solution near the horizon, ($r \rightarrow 0$). There $\mathcal{V}(Y)$ can be approximated as follows.

$$\mathcal{V}_{hor}(Y) = \frac{1}{3} (Y^I H_I)_{hor} = \frac{1}{3} (Y^I)_{hor} \left(\frac{q_I}{r^2} \right). \quad (17)$$

However, $Y_{hor}^I = \mathcal{V}_{hor}^{1/3}(Y) X_{hor}^I$, and thus

$$\mathcal{V}_{hor}^{2/3}(Y) = \frac{1}{3} \frac{(Z_e)_{cr}}{r^2} \quad (18)$$

where $Z_e = q_I X^I$ is the electric central charge, and $(Z_e)_{cr}$ is its value at the horizon. Moreover, equation (10) which defines the moduli over space-time becomes near the horizon

$$(Z_e X_I)_{hor} = q_I, \quad (19)$$

which is what one obtains from the extremisation of the central charge. Comparison of the form of our metric near the horizon with (16), implies that the two forms are related by

$$r_0^2 = \frac{(Z_e)_{cr}}{3}, \quad \hat{r} = \frac{r^2}{2r_0}. \quad (20)$$

The proof for the enhancement of supersymmetry for the double extreme black hole solution therefore remains unchanged for the general electric BPS solution as the metrics near the horizon are identical.

Here we complete the analysis by solving for the Killing spinor near the horizon. There the gravitino supersymmetry transformations become

$$\begin{aligned}
\delta\psi_t &= \nabla_t \epsilon = \left(\partial_t + ir^2 \left(\frac{3}{(Z_e)_{\text{cr}}} \right)^{\frac{3}{2}} \Gamma_1 (1 + i\Gamma_0) \right) \epsilon, \\
\delta\psi_r &= \nabla_r \epsilon = \left(\partial_r + \frac{i}{r} \Gamma_0 \right) \epsilon, \\
\delta\psi_\theta &= \nabla_\theta \epsilon = \left(\partial_\theta + \frac{i}{2} \Gamma_{012} \right) \epsilon, \\
\delta\psi_\phi &= \nabla_\phi \epsilon = \left(\partial_\phi - \frac{1}{2} \cos \theta \Gamma_{23} + \frac{i}{2} \Gamma_{013} \sin \theta \right) \epsilon, \\
\delta\psi_\psi &= \nabla_\psi \epsilon = \left(\partial_\psi + \frac{1}{2} \sin \theta \Gamma_{24} + \frac{i}{2} \Gamma_{014} \cos \theta \right) \epsilon,
\end{aligned} \tag{21}$$

where we have used that near the horizon $X^I F_{tr}^I = -\frac{6r}{(Z_e)_{\text{cr}}}$.

The integrability conditions of the gravitino transformations impose no conditions on the Killing spinor ϵ , which implies that one has full supersymmetry. Setting all the variations in (21) to zero can be solved by the Killing spinor

$$\epsilon(r, \theta, \phi, \psi) = e^{-\frac{i}{2} \Gamma_{012} \theta} e^{\frac{1}{2} \Gamma_{23} \phi} e^{-\frac{i}{2} \Gamma_{014} \psi} \left(\frac{1}{2r} - i \text{tr} \left(\frac{3}{(Z_e)_{\text{cr}}} \right)^{\frac{3}{2}} \Gamma_1 (1 + i\Gamma_0) \right) \varepsilon_0, \tag{22}$$

where ε_0 is a constant spinor.

3 BPS Magnetic Solutions

In this section we construct general BPS magnetic string-like solutions which break half of supersymmetry of $d = 5$, $N = 2$ supergravity theory coupled to vector supermultiplets. As a general metric for the magnetic string-solution we write

$$ds^2 = e^{2V} (-dt^2 + dz^2) + e^{2U} (d\vec{x})^2, \tag{23}$$

where $(d\vec{x})^2 = (dx^1)^2 + (dx^2)^2 + (dx^3)^2$, and (x^1, x^2, x^3) are the transverse dimensions. The functions U and V are taken to be independent of (t, z) . The non-vanishing Fünfbins for the metric in (23) are

$$e_t^0 = e_z^1 = e^V, \quad e_m^a = e^U \delta_m^a. \tag{24}$$

For the spin connections one obtains

$$\begin{aligned}
\omega_{ta0} &= \partial_m V e^{V-U} \delta_a^m, \\
\omega_{za1} &= -\partial_m V e^{V-U} \delta_a^m, \\
\omega_{pab} &= \partial_n U (\delta^{an} \delta_p^b - \delta^{nb} \delta_p^a).
\end{aligned} \tag{25}$$

First we consider the supersymmetry variation of the gauginos and find the conditions that must be satisfied by a supersymmetric bosonic configuration. These conditions are obtained by setting the fermions and their supersymmetry variations to zero. Using (4), the gaugino transformation can be rewritten in the form

$$\delta\lambda_i = -\frac{1}{4} \left(G_{IJ} \partial_i X^I \Gamma^{\mu\nu} F_{\mu\nu}^J - 3i \Gamma^\mu \partial_\mu X_I \partial_i X^I \right) \epsilon. \quad (26)$$

The vanishing of the gaugino transformation thus gives

$$G_{IJ} \partial_i X^I \left(\Gamma^{mn} F_{mn}^J + 2i \partial_m X^J \Gamma^m \right) \epsilon = 0, \quad (27)$$

where we have used the relation

$$G_{IJ} \partial_\mu X^J = -\frac{3}{2} \partial_\mu X_I, \quad (28)$$

which follows from very special geometry. Assuming that the susy parameter satisfies

$$\epsilon = \Gamma_1 \Gamma_0 \epsilon, \quad (29)$$

a condition which implies that half of supersymmetry is broken. Using (29), it can be shown that

$$\Gamma^{mn} \epsilon = i e^{-3U} \epsilon^{mnp} \Gamma_p \epsilon, \quad (30)$$

If one also writes $X^I = e^{-U} H^I$, then (27) together with the relation $X^I \partial_i X^I = 0$, implies the following

$$\Gamma^{mn} F_{mn}^I \epsilon = -2i e^{-U} \partial_p H^I \Gamma^p \epsilon. \quad (31)$$

This, with (30) fix the gauge field strengths to be

$$F_{mn}^I = -\epsilon_{mnp} \partial_p H^I. \quad (32)$$

Moreover, using the relation $\frac{1}{6} C_{IJK} X^I X^J X^K = 1$, one concludes that

$$e^{3U} = \frac{1}{6} C_{IJK} H^I H^J H^K. \quad (33)$$

Therefore, using the relations of very special geometry, the vanishing of the gaugino supersymmetry variation for a choice of the supersymmetry parameter, fixes the relation between the gauge field strengths, X^I and U .

The vanishing of the gravitino supersymmetry transformation fixes the metric as well as the supersymmetry parameter itself. The vanishing of the t -component gives

$$\delta\psi_t = \left(\partial_t + \frac{1}{4} \omega_t^{ab} \Gamma_{ab} + \frac{i}{8} \Gamma_t^{mn} X_I F_{mn}^I \right) \epsilon = 0. \quad (34)$$

Using our Ansatz, we have

$$\frac{1}{4} \omega_t^{ab} \Gamma_{ab} \epsilon = \frac{1}{2} \omega_t^{a0} \Gamma_{a0} \epsilon = -\frac{1}{2} \partial_m V e^{V-U} \delta_m^a \Gamma_a \Gamma_0 \epsilon. \quad (35)$$

Moreover, using the expressions for the scalar fields and gauge field strengths we get

$$\frac{i}{8}X_I\Gamma_t^{mn}F_{mn}^I = \frac{1}{4}e^{V-U}\partial_m U\delta_a^m\Gamma_{a0}\epsilon. \quad (36)$$

Therefore, from (34), we get the conditions

$$V = -\frac{1}{2}U, \quad \partial_t\epsilon = 0. \quad (37)$$

Similarly, the z component of the gravitino supersymmetry variation gives

$$\delta\psi_z = \left(\partial_z + \frac{1}{4}\omega_z^{ab}\Gamma_{ab} + \frac{i}{8}X_I\Gamma_z^{mn}F_{mn}^I\right)\epsilon. \quad (38)$$

Using $X_I\Gamma^{mn}F_{mn}^I\epsilon = -2i\Gamma^p\partial_p U\epsilon$, $\Gamma_z = e_z^1\Gamma_1$, $\Gamma^p = e_a^p\Gamma^a$, we deduce that $\partial_z\epsilon = 0$. Moreover, one obtains from the vanishing of the space-components of the gravitino supersymmetry variations

$$(\partial_m + \frac{1}{4}\partial_m U)\epsilon = 0. \quad (39)$$

Thus the Killing spinor is given by

$$\epsilon = e^{-\frac{1}{4}U}\epsilon_0, \quad (40)$$

where ϵ_0 is a constant spinor satisfying $\epsilon = \Gamma_1\Gamma_0\epsilon$.

Here we give a summary of what we have done so far. We are seeking a background which breaks supersymmetry and we have obtained the following configuration

$$F_{mn}^I = -\epsilon_{mnp}\partial_p H^I, \quad X^I = e^{-U}H^I, \quad V = -\frac{1}{2}U. \quad (41)$$

In order to fix the scalar functions H^I , one has to solve for the equations of motion and the Bianchi identities for the gauge fields. Doing so, the H^I are then fixed and are given by a set of harmonic functions

$$H^I = g^I + \frac{p^I}{r}, \quad (42)$$

where g^I are constants and p^I are magnetic charges. As for the electric black hole solutions, we define new coordinates $Y^I = e^U X^I$, in terms of which the magnetic string solution takes the following simple form

$$\begin{aligned} ds^2 &= \mathcal{V}(Y)^{-\frac{1}{3}}(-dt^2 + dz^2) + \mathcal{V}(Y)^{\frac{2}{3}}(dr^2 + r^2 d\theta^2 + r^2 \sin^2\theta d\phi^2), \\ F_{mn}^I &= -\epsilon_{mnp}\partial_p H^I, \\ Y^I &= H^I. \end{aligned} \quad (43)$$

In the electric case, the horizon geometry is given by $AdS_2 \times S^3$ and the black hole entropy, related to the horizon volume S^3 , is given in terms of the extremised electric

central charge. For the magnetically charged $d = 5$ BPS black string, one similarly find that for our solution the extremised value of the BPS tension is related to the volume of the S^2 where the near horizon geometry is given by $AdS_3 \times S^2$. This can be easily demonstrated. The magnetic central charge related to the tension of the magnetic string states is defined by $Z_m = p^I X_I$. To find the extremum we set

$$\partial_i Z_m = \partial_i (X_I) p^I = \frac{1}{3} C_{IJK} X^I \partial_i (X^J) p^K = 0. \quad (44)$$

It follows that the critical values of X^I and its dual are given by [18]

$$X^I = \frac{p^I}{Z_m}, \quad X_I = \frac{1}{6} \frac{C_{IJK} p^J p^K}{Z_m^2} \quad (45)$$

and thus the critical value of the BPS string tension is

$$(Z_m)_{\text{cr}}^3 = \frac{1}{6} C_{IJK} p^I p^J p^K. \quad (46)$$

At the near horizon ($r \rightarrow 0$) we have

$$\begin{aligned} e^{3U} &= \frac{1}{6} C_{IJK} H^I H^J H^K \rightarrow \frac{1}{6r^3} C_{IJK} p^I p^J p^K \equiv \frac{(Z_m)_{\text{cr}}^3}{r^3} \\ X^I &= e^{-U} H^I \rightarrow \frac{p^I}{(Z_m)_{\text{cr}}} \\ X_I &= \frac{1}{6} C_{IJK} H^J H^K \rightarrow \frac{1}{6} C_{IJK} \frac{p^J p^K}{(Z_m)_{\text{cr}}^2}. \end{aligned} \quad (47)$$

Therefore the values of the moduli at the horizon are those which extremise the magnetic central charge. With these limits, the metric and the gauge fields at the horizon could then be expressed in the form

$$\begin{aligned} ds^2 &= \frac{r}{(Z_m)_{\text{cr}}} (-dt^2 + dz^2) + \frac{(Z_m)_{\text{cr}}^2}{r^2} dr^2 + (Z_m)_{\text{cr}}^2 (d\theta^2 + \sin^2 \theta d\phi^2), \\ F_{\theta\phi}^I &= q^I \sin \theta. \end{aligned} \quad (48)$$

We now demonstrate the enhancement of supersymmetry at the horizon for the magnetic solutions. For the near horizon metric, the gaugino transformation reduces to

$$\delta\lambda_i = \left(\frac{1}{3(Z_m)_{\text{cr}}^2} \partial_i (q^I X_I) \Gamma_{34} + \frac{3i}{4(Z_m)_{\text{cr}}} r (\partial_r X_I \partial_i X^I) \Gamma_2 \right) \epsilon \quad (49)$$

The above variation vanishes identically without any conditions imposed on the supersymmetry parameter. This is basically due to the fact that the magnetic central charge is

extremised at the horizon. From the gravitino supersymmetry transformations near the horizon we get the equations

$$\begin{aligned}
\delta\psi_t &= \nabla_t \epsilon = \left(\partial_t + \frac{r^{\frac{1}{2}}}{4(Z_m)_{\text{cr}}^{\frac{3}{2}}} (\Gamma_0 - \Gamma_1) \Gamma_2 \right) \epsilon, \\
\delta\psi_z &= \nabla_z \epsilon = \left(\partial_z - \frac{r^{\frac{1}{2}}}{4(Z_m)_{\text{cr}}^{\frac{3}{2}}} (\Gamma_0 - \Gamma_1) \Gamma_2 \right) \epsilon, \\
\delta\psi_r &= \nabla_r \epsilon = \left(\partial_r + \frac{1}{4r} \Gamma_0 \Gamma_1 \right) \epsilon, \\
\delta\psi_\theta &= \nabla_\theta \epsilon = \left(\partial_\theta - \frac{i}{2} \Gamma_4 \right) \epsilon, \\
\delta\psi_\phi &= \nabla_\phi \epsilon = \left(\partial_\phi + \frac{i}{2} \sin \theta \Gamma_3 - \frac{1}{2} \cos \theta \Gamma_3 \Gamma_4 \right) \epsilon,
\end{aligned} \tag{50}$$

The integrability conditions

$$[\nabla_\mu, \nabla_\nu] \epsilon = 0 \tag{51}$$

can be seen to be satisfied with no restrictions on the Killing spinor ϵ . This implies that supersymmetry is fully restored. The solution of the Killing spinor is given by ⁶

$$\epsilon(t, z, r, \theta, \phi) = e^{\frac{i}{2} \Gamma_4 \theta} e^{\frac{1}{2} \Gamma_3 \Gamma_4 \phi} \left((1 + \Gamma_0 \Gamma_1) r^{\frac{-1}{4}} - \frac{(t - z)}{2(Z_m)_{\text{cr}}^{\frac{3}{2}}} r^{\frac{1}{4}} (\Gamma_0 - \Gamma_1) \Gamma_2 \right) \kappa_0. \tag{52}$$

where κ_0 is a constant spinor. The structure of the horizon is $AdS_3 \times S^2$ with curvatures

$$\begin{aligned}
R_{ab} &= -\frac{1}{4(Z_m)_{\text{cr}}^2} \eta'_{ab}, & \eta'_{ab} &= (-1, 1, 1), & a, b &= 0, 1, 2 \\
R_{\alpha\beta} &= \frac{1}{2(Z_m)_{\text{cr}}^2} \delta_{\alpha\beta}, & \alpha, \beta &= 3, 4.
\end{aligned} \tag{53}$$

In conclusion, with the help of the relations of very special geometry, BPS electric and magnetic solutions of M-theory on a Calabi-Yau manifold were obtained. Our analysis applies to all $N = 2$ five dimensional supergravity and not only to models with higher-dimensional origin, *i.e.*, those obtained from compactifying 11-dimensional supergravity on a Calabi-Yau manifold. The solutions depend on the prepotential which defines the theory. We note that in the electric solutions, the rescaled four-cycles Y_I are given in terms of harmonic functions and thus in order to find the explicit space-time solution one has to solve for the two-cycles X^I . In the magnetic case, however, it is the rescaled two-cycle which is given in terms of a harmonic function and thus the explicit space-time solution is known for all prepotentials. We have demonstrated that supersymmetry is enhanced at the near horizon of the general electric and magnetic solutions. The solutions for the Killing spinors of the near horizon geometries, $AdS_2 \times S^3$ and $AdS_3 \times S^2$ are given.

⁶the dependence of the Killing spinor on Z_m can be absorbed by the redefinition of coordinates, $r = r'(Z_m)_{\text{cr}}^3$

References

- [1] S. Ferrara, R. Kallosh and A. Strominger, *Phys. Rev.* **D52** (1995) 5412; S. Ferrara and R. Kallosh, *Phys. Rev.* **D54** (1996) 1514; S. Ferrara, R. Kallosh, *Phys. Rev.* **D54** (1996) 1525.
- [2] D. Lüst, hep-th/9803072; D. Youm, hep-th/9710046; A. W. Peet, *Class. Quant. Grav.* **15** (1998) 3291.
- [3] B. Craps, F. Roose, W. Troost, A Van Proeyen, *Nucl. Phys.* **B503** (1997) 565.
- [4] B. de Wit and A. Van Proeyen, *Phys. Lett.* **293** (1992) 94.
- [5] W. A. Sabra, *Mod. Phys. Lett.* **A 12** (1997) 2585; *Nucl. Phys.* **B 510** (1998) 247; K. Behrndt, D. Lüst and W. A. Sabra, *Nucl. Phys.* **B510** (1998) 264.
- [6] J. B. Hartle and S. W. Hawking , *Commun. Math. Phys.* **26** (1972) 87.
- [7] W. A. Sabra, *Mod. Phys. Lett.* **A 13** (1998) 239 ; A. Chamseddine and W. A. Sabra, *Phys. Lett.* **B426** (1998) 36.
- [8] G.W. Gibbons, *Nucl. Phys.* **B207** (1982) 337.
- [9] R. Kallosh and A. Peet, *Phys. Rev.* **D46** (1992) 5223.
- [10] G. W. Gibbons and P. K. Townsend, *Phys. Rev. Lett.* **71**, (1993) 3754.
- [11] R. Kallosh, M. Shmakova and W. K. Wong, *Phys. Rev.* **D54** (1996) 6284.
- [12] A. Chamseddine, S. Ferrara, G.W. Gibbons and R. Kallosh, *Phys. Rev.* **D 55** (1997) 3647.
- [13] R. Kallosh, A. Rajaraman, W. K. Wong, *Phys. Rev.* **D55** (1997) 3249.
- [14] G. W. Gibbons, D. Kastor, L. A. J. London, P. K. Townsend and J. Traschen, *Nucl. Phys.* **B416** (1994) 850.
- [15] K. Behrndt, *Phys. Lett.* **B396** (1997) 77.
- [16] M. Günaydin, G. Sierra, and P. K. Townsend, *Class. Quant. Grav* **3** (1986) 763.
- [17] M. Günaydin, G. Sierra, and P. K. Townsend, *Nucl. Phys.* **B242** (1984) 244; *Nucl. Phys.* **B253** (1985) 573.
- [18] A. Chou, R. Kallosh, J. Rahmfeld, S.-J. Rey, M. Shmakova, W. K. Wong, *Nucl. Phys.* **B508** (1997) 147